

3 5 IMPORTANT REASONS

why you and your child's doctor
may consider **LUPRON DEPOT-PED®**

**YOU CAN'T CONTROL
Everything they DO
But you can help control
Central Precocious Puberty (CPP)
with LUPRON DEPOT-PED**

USE FOR LUPRON DEPOT-PED® (leuprolide acetate for depot suspension)


LUPRON DEPOT-PED 7.5 mg, 11.25 mg, and 15 mg for 1-month, 11.25 mg and 30 mg for 3-month, and 45 mg for 6-month administration are prescribed for the treatment of children with central precocious puberty (CPP). It is not known if LUPRON DEPOT-PED is safe and effective in children less than 1 year old.

SAFETY CONSIDERATIONS

- Your child should not receive LUPRON DEPOT-PED if she/he is allergic to any of the ingredients or she is or may become pregnant.
- Increased signs and symptoms of puberty during the first 2 to 4 weeks of treatment may occur. Contact your child's doctor if signs of puberty continue after the second month of treatment.

Please see additional Important Safety Information on pages 6-7.

Click [here](#) for Consumer Brief Summary or visit <https://www.rxabbvie.com/pdf/lupronpediatric.pdf> for full Prescribing Information.




Reason 1:

#1 prescribed treatment for CPP*



Reason 2:

ONLY CPP treatment with over 30 years of proven efficacy and safety[†]



Treating CPP since 1993!

*Data sourced as of September 2025.

[†]FDA approved in 1993.

SAFETY CONSIDERATIONS (continued)

- Some people taking gonadotropin-releasing hormone (GnRH) agonists, like LUPRON DEPOT-PED, have had new or worsened mental (psychiatric) problems. Mental (psychiatric) problems may include emotional symptoms such as crying, irritability, restlessness (impatience), anger, and acting aggressive. Call your child's doctor right away if your child has any new or worsening mental symptoms or problems while taking LUPRON DEPOT-PED.

Please see additional Important Safety Information on pages 6–7.

Click [here](#) for **Consumer Brief Summary** or visit
<https://www.rxabbvie.com/pdf/lupronpediatric.pdf> for
full **Prescribing Information**.

LUPRON DEPOT-PED®
(leuprolide acetate for depot suspension)

Reason 3:

ONLY CPP treatment with an 18-year, long-term efficacy and safety study

In a study of 1-month dosing,* LUPRON DEPOT-PED provided control by **pausing early puberty within 4 weeks**. While on treatment, your child's physical signs of early puberty should be paused. Genital and breast development should stop or reverse.

After treatment with 1-month LUPRON DEPOT-PED, the reproductive development of 20 patients aged 18-26 was evaluated:



80% (16 of 20) reported normal menstrual cycles



12 pregnancies in 7 patients (including multiple pregnancies in 4 patients)



4 of 4 women who tried to become pregnant did

On average, children on therapy grew taller than the height predicted for them when they were first diagnosed with CPP.



*The study included 55 children (49 females and 6 males) with CPP treated with LUPRON DEPOT-PED for an average of 3.9 years who were then followed after treatment for an average of 3.5 years.

SAFETY CONSIDERATIONS (continued)

- Seizures have been observed in patients taking GnRH agonists, like LUPRON DEPOT-PED, and the risk of seizure may be higher in patients with or without a history of seizures, epilepsy, brain or brain vessel (cerebrovascular) problems or tumors, and in patients taking medications that have been connected to seizures, such as bupropion or selective serotonin reuptake inhibitors (SSRIs). Call your child's doctor right away if your child has a seizure while taking LUPRON DEPOT-PED.

Please see additional Important Safety Information on pages 6-7.

Click [here](#) for **Consumer Brief Summary** or visit
<https://www.rxabbvie.com/pdf/lupronpediatric.pdf> for
full **Prescribing Information**.


Reason 4:

Individualized dosing: One size doesn't fit all

The ONLY CPP treatment with multiple dosing options to meet your child's needs

Dosing options available:

- **Every 6 months**
- **Every 3 months**
- **Every month**



Reason 5:

The thinnest needle for CPP injections*



Injection site reactions including sterile abscess, injection site pain, and injection site swelling* occurred in $\geq 2\%$ of patients with CPP in clinical trials.

**LUPRON DEPOT-PED
supports you every
step of the way**

Visit LupronPed.com/LupronSupport-Plus
for information on insurance coverage,
helpful resources, and to see
if you are eligible for our
\$10 savings card

*Most events were mild or moderate in severity.

SAFETY CONSIDERATIONS (continued)

- Severe cutaneous (skin) adverse reactions may happen during treatment with GnRH agonists like LUPRON DEPOT-PED. Stop LUPRON DEPOT-PED and call your child's doctor right away if your child has a skin rash or acne; dry skin; itching; blisters on their skin; redness or swelling of their face, hands, or soles of their feet; blisters or sores in their mouth; peeling of their skin; fever; muscle or joint aches; or swollen glands.

Please see additional Important Safety Information on pages 6–7.

Click [here](#) for **Consumer Brief Summary** or visit
<https://www.rxabbvie.com/pdf/lupronpediatric.pdf> for
full **Prescribing Information**.

LUPRON DEPOT-PED®
(leuprolide acetate for depot suspension)

5 IMPORTANT REASONS

why you and your child's doctor
may consider **LUPRON DEPOT-PED®**

- **#1 prescribed** treatment for CPP*
- Over **30 years** of **proven** efficacy and **safety**†
- **18-year**, long-term efficacy and **safety study**
- **Individualized** dosing
- The **thinnest needle** for CPP injections



Scan this QR code or visit

LupronPed.com

for more information

*Data sourced as of September 2025.

†FDA approved in 1993.




SAFETY CONSIDERATIONS (continued)

- Increased pressure in the fluid around the brain can happen in children taking GnRH agonist medicines, including LUPRON DEPOT-PED. Call your child's doctor right away if your child has any of the following symptoms during treatment with LUPRON DEPOT-PED: headache, eye problems (including blurred vision, double vision, and decreased eyesight), eye pain, ringing in the ears, dizziness, or nausea.
- Your child's pubertal development could begin again if the injection schedule is not followed.

Please see additional Important Safety Information on pages 6–7.

Click [here](#) for **Consumer Brief Summary** or visit

<https://www.rxabbvie.com/pdf/lupronpediatric.pdf> for
full **Prescribing Information**.



USE AND IMPORTANT SAFETY INFORMATION

Use for LUPRON DEPOT-PED®

(leuprolide acetate for depot suspension)

LUPRON DEPOT-PED 7.5 mg, 11.25 mg, and 15 mg for 1-month, 11.25 mg and 30 mg for 3-month, and 45 mg for 6-month administration are prescribed for the treatment of children with central precocious puberty (CPP).

It is not known if LUPRON DEPOT-PED is safe and effective in children less than 1 year old.

Important Safety Information for LUPRON DEPOT-PED

What is the most important information I should know about LUPRON DEPOT-PED?

- **During the first 2 to 4 weeks of treatment, LUPRON DEPOT-PED can cause an increase in some hormones.** During this time, you may notice more signs of puberty in your child, including vaginal bleeding. **Call your child's doctor if these signs continue after the second month of treatment with LUPRON DEPOT-PED.**

- **Some people taking gonadotropin-releasing hormone (GnRH) agonists like LUPRON DEPOT-PED have had new or worsened mental (psychiatric) problems.** Mental (psychiatric) problems may include emotional symptoms such as:

- Crying
- Irritability
- Restlessness (impatience)
- Anger
- Acting aggressive

Call your child's doctor right away if your child has any new or worsening mental symptoms or problems while taking LUPRON DEPOT-PED.

- **Some people taking GnRH agonists like LUPRON DEPOT-PED have had seizures. The risk of seizures may be higher in people who:**

- Have a history of seizures
- Have a history of epilepsy
- Have a history of brain or brain vessel (cerebrovascular) problems or tumors
- Are taking a medicine that has been connected to seizures, such as bupropion or selective serotonin reuptake inhibitors (SSRIs)

Please see additional Important Safety Information on page 7.

Click [here](#) for **Consumer Brief Summary** or visit <https://www.rxabbvie.com/pdf/lupronpediatric.pdf> for full **Prescribing Information**.

Seizures have also happened in people who have not had any of these problems. **Call your child's doctor right away if your child has a seizure while taking LUPRON DEPOT-PED.**

- **Severe cutaneous (skin) adverse reactions may happen during treatment with GnRH agonists like LUPRON DEPOT-PED. Stop LUPRON DEPOT-PED and call your child's doctor right away if your child has any of the following signs or symptoms during treatment with LUPRON DEPOT-PED:**

- Skin rash or acne
- Dry skin
- Itching
- Blisters on your skin
- Redness or swelling of your face, hands, or soles of your feet
- Blisters or sores in your mouth
- Peeling of your skin
- Fever
- Muscle or joint aches
- Swollen glands

- **Increased pressure in the fluid around the brain can happen in children taking gonadotropin-releasing hormone (GnRH) agonist medicines, including LUPRON DEPOT-PED. Call your child's doctor right away if your child has any of the following symptoms during treatment with LUPRON DEPOT-PED:**

- Headache
- Eye problems, including blurred vision, double vision, and decreased eyesight
- Eye pain
- Ringing in the ears
- Dizziness
- Nausea

LUPRON DEPOT-PED should not be taken if your child is:

- Allergic to GnRH, GnRH agonist medicines, or any ingredients in LUPRON DEPOT-PED. See the end of the Medication Guide for a complete list of ingredients in LUPRON DEPOT-PED.
- Pregnant or becomes pregnant. LUPRON DEPOT-PED can cause birth defects or loss of the baby. If your child becomes pregnant, call your child's doctor.

(continued on page 7)

LUPRON DEPOT-PED®
(leuprolide acetate for depot suspension)

USE AND IMPORTANT SAFETY INFORMATION (CONTINUED)

Important Safety Information for LUPRON DEPOT-PED (continued)

Before your child receives LUPRON DEPOT-PED, tell their doctor about all of your child's medical conditions, including if they:

- Have a history of mental (psychiatric) problems
- Have a history of seizures
- Have a history of epilepsy
- Have a history of brain or brain vessel (cerebrovascular) problems or tumors
- Are taking a medicine that has been connected to seizures, such as bupropion or selective serotonin reuptake inhibitors (SSRIs)
- Are breastfeeding or plan to breastfeed. It is not known if LUPRON DEPOT-PED passes into the breast milk

Tell your child's doctor about all the medicines your child takes, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

How will your child receive LUPRON DEPOT-PED?

- Your child's doctor should do tests to make sure your child has CPP before treating them with LUPRON DEPOT-PED.
- LUPRON DEPOT-PED is given as a single-dose injection into your child's muscle each month, every 3 months, or every 6 months by a doctor or trained nurse. Your child's doctor will decide how often your child will receive the injection.
- Keep all scheduled visits to the doctor. If a scheduled dose is missed, your child may start having signs of puberty again. The doctor will do regular exams and blood tests to check for signs of puberty.

What are the possible side effects of LUPRON DEPOT-PED?

LUPRON DEPOT-PED may cause serious side effects. See "What is the most important information I should know about LUPRON DEPOT-PED?"

• The most common side effects of LUPRON DEPOT-PED received 1 time each month include:

- Injection site reactions such as pain, swelling, and abscess
- Weight gain
- Pain throughout body

- Headache
- Acne or red, itchy rash and white scales (seborrhea)
- Serious skin rash (erythema multiforme)
- Mood changes
- Swelling of vagina (vaginitis), vaginal bleeding, and vaginal discharge

• The most common side effects of LUPRON DEPOT-PED received every 3 months include:

- Injection site reactions such as pain and swelling
- Weight gain
- Headache
- Mood changes

• The most common side effects of LUPRON DEPOT-PED received every 6 months include:

- Injection site reactions such as pain and swelling
- Headache
- Mood changes
- Upper stomach pain
- Diarrhea
- Bleeding
- Nausea and vomiting
- Fever
- Itching
- Pain in extremities
- Rash
- Back pain
- Ligament sprain
- Weight gain
- Fracture
- Breast tenderness
- Difficulty sleeping
- Chest pain
- Excessive sweating

These are not all the possible side effects of LUPRON DEPOT-PED. Call your child's doctor for medical advice about side effects.

This is the most important information to know about LUPRON DEPOT-PED. For more information, talk to your child's doctor or healthcare provider.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088.

If you are having difficulty paying for your child's medicine, AbbVie may be able to help. Visit AbbVie.com/PatientAccessSupport to learn more.

Please see additional Important Safety Information on page 6.

Click [here](#) for **Consumer Brief Summary** or visit <https://www.rxabbvie.com/pdf/lupronpediatric.pdf> for full **Prescribing Information**.